


**4.37** Two infinite lines of charge, both parallel to the  $z$ -axis, lie in the  $x$ - $z$  plane, one with density  $\rho_\ell$  and located at  $x = a$  and the other with density  $-\rho_\ell$  and located at  $x = -a$ . Obtain an expression for the electric potential  $V(x, y)$  at a point  $P = (x, y)$  relative to the potential at the origin.



**Solution:** According to the result of Problem 4.33, the electric potential difference between a point at a distance  $r_1$  and another at a distance  $r_2$  from a line charge of density  $\rho_l$  is

$$V = \frac{\rho_l}{2\pi\epsilon_0} \ln\left(\frac{r_2}{r_1}\right).$$

Applying this result to the line charge at  $x = a$ , which is at a distance  $a$  from the origin:

$$\begin{aligned} V' &= \frac{\rho_l}{2\pi\epsilon_0} \ln\left(\frac{a}{r'}\right) \quad (r_2 = a \text{ and } r_1 = r') \\ &= \frac{\rho_l}{2\pi\epsilon_0} \ln\left(\frac{a}{\sqrt{(x-a)^2 + y^2}}\right). \end{aligned}$$

Similarly, for the negative line charge at  $x = -a$ ,

$$\begin{aligned} V'' &= \frac{-\rho_l}{2\pi\epsilon_0} \ln\left(\frac{a}{r''}\right) \quad (r_2 = a \text{ and } r_1 = r') \\ &= \frac{-\rho_l}{2\pi\epsilon_0} \ln\left(\frac{a}{\sqrt{(x+a)^2 + y^2}}\right). \end{aligned}$$

The potential due to both lines is

$$V = V' + V'' = \frac{\rho_l}{2\pi\epsilon_0} \left[ \ln\left(\frac{a}{\sqrt{(x-a)^2 + y^2}}\right) - \ln\left(\frac{a}{\sqrt{(x+a)^2 + y^2}}\right) \right].$$

At the origin,  $V = 0$ , as it should be since the origin is the reference point. The potential is also zero along all points on the  $y$ -axis ( $x = 0$ ).